

# Chapter Four

# MOTION IN A PLANE

## MCQ I

- 4.1** The angle between  $\mathbf{A} = \hat{\mathbf{i}} + \hat{\mathbf{j}}$  and  $\mathbf{B} = \hat{\mathbf{i}} - \hat{\mathbf{j}}$  is  
(a)  $45^\circ$  (b)  $90^\circ$  (c)  $-45^\circ$  (d)  $180^\circ$
- 4.2** Which one of the following statements is true?  
(a) A scalar quantity is the one that is conserved in a process.  
(b) A scalar quantity is the one that can never take negative values.  
(c) A scalar quantity is the one that does not vary from one point to another in space.  
(d) A scalar quantity has the same value for observers with different orientations of the axes.
- 4.3** Figure 4.1 shows the orientation of two vectors  $\mathbf{u}$  and  $\mathbf{v}$  in the XY plane.

If  $\mathbf{u} = a\hat{\mathbf{i}} + b\hat{\mathbf{j}}$  and

$$\mathbf{v} = p\hat{\mathbf{i}} + q\hat{\mathbf{j}}$$




Fig. 4.1

which of the following is correct?

- (a)  $a$  and  $p$  are positive while  $b$  and  $q$  are negative.
- (b)  $a$ ,  $p$  and  $b$  are positive while  $q$  is negative.
- (c)  $a$ ,  $q$  and  $b$  are positive while  $p$  is negative.
- (d)  $a$ ,  $b$ ,  $p$  and  $q$  are all positive.

**4.4** The component of a vector  $\mathbf{r}$  along  $X$ -axis will have maximum value if

- (a)  $\mathbf{r}$  is along positive  $Y$ -axis
- (b)  $\mathbf{r}$  is along positive  $X$ -axis
- (c)  $\mathbf{r}$  makes an angle of  $45^\circ$  with the  $X$ -axis
- (d)  $\mathbf{r}$  is along negative  $Y$ -axis

**4.5** The horizontal range of a projectile fired at an angle of  $15^\circ$  is 50 m. If it is fired with the same speed at an angle of  $45^\circ$ , its range will be

- (a) 60 m
- (b) 71 m
- (c) 100 m
- (d) 141 m

**4.6** Consider the quantities, pressure, power, energy, impulse, gravitational potential, electrical charge, temperature, area. Out of these, the only vector quantities are

- (a) Impulse, pressure and area
- (b) Impulse and area
- (c) Area and gravitational potential
- (d) Impulse and pressure

**4.7** In a two dimensional motion, instantaneous speed  $v_0$  is a positive constant. Then which of the following are necessarily true?

- (a) The average velocity is not zero at any time.
- (b) Average acceleration must always vanish.
- (c) Displacements in equal time intervals are equal.
- (d) Equal path lengths are traversed in equal intervals.

**4.8** In a two dimensional motion, instantaneous speed  $v_0$  is a positive constant. Then which of the following are necessarily true?

- (a) The acceleration of the particle is zero.
- (b) The acceleration of the particle is bounded.
- (c) The acceleration of the particle is necessarily in the plane of motion.
- (d) The particle must be undergoing a uniform circular motion

**4.9** Three vectors  $\mathbf{A}, \mathbf{B}$  and  $\mathbf{C}$  add up to zero. Find which is false.

- (a)  $(\mathbf{A} \times \mathbf{B}) \times \mathbf{C}$  is not zero unless  $\mathbf{B}, \mathbf{C}$  are parallel
- (b)  $(\mathbf{A} \times \mathbf{B}) \cdot \mathbf{C}$  is not zero unless  $\mathbf{B}, \mathbf{C}$  are parallel
- (c) If  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  define a plane,  $(\mathbf{A} \times \mathbf{B}) \times \mathbf{C}$  is in that plane
- (d)  $(\mathbf{A} \times \mathbf{B}) \cdot \mathbf{C} = |\mathbf{A}| |\mathbf{B}| |\mathbf{C}| \cos C^2 = A^2 + B^2$

**4.10** It is found that  $|\mathbf{A} + \mathbf{B}| = |\mathbf{A}|$ . This necessarily implies,

- (a)  $\mathbf{B} = \mathbf{0}$
- (b)  $\mathbf{A}, \mathbf{B}$  are antiparallel
- (c)  $\mathbf{A}, \mathbf{B}$  are perpendicular
- (d)  $\mathbf{A} \cdot \mathbf{B} \leq 0$

## MCQ II

**4.11** Two particles are projected in air with speed  $v_o$  at angles  $\theta_1$  and  $\theta_2$  (both acute) to the horizontal, respectively. If the height reached by the first particle is greater than that of the second, then tick the right choices

- (a) angle of projection :  $q_1 > q_2$
- (b) time of flight :  $T_1 > T_2$
- (c) horizontal range :  $R_1 > R_2$
- (d) total energy :  $U_1 > U_2$ .

**4.12** A particle slides down a frictionless parabolic ( $y = x^2$ ) track (A – B – C) starting from rest at point A (Fig. 4.2). Point B is at the vertex of parabola and point C is at a height less than that of point A. After C, the particle moves freely in air as a projectile. If the particle reaches highest point at P, then

- (a) KE at P = KE at B
- (b) height at P = height at A
- (c) total energy at P = total energy at A
- (d) time of travel from A to B = time of travel from B to P.

**4.13** Following are four different relations about displacement, velocity and acceleration for the motion of a particle in general. Choose the incorrect one (s) :

(a)  $\mathbf{v}_{av} = \frac{1}{2} [\mathbf{v}(t_1) + \mathbf{v}(t_2)]$

(b)  $\mathbf{v}_{av} = \frac{\mathbf{r}(t_2) - \mathbf{r}(t_1)}{t_2 - t_1}$




Fig. 4.2

(c)  $\mathbf{r} = \frac{1}{2}(\mathbf{v}(t_2) - \mathbf{v}(t_1))(t_2 - t_1)$

(d)  $\mathbf{a}_{av} = \frac{\mathbf{v}(t_2) - \mathbf{v}(t_1)}{t_2 - t_1}$

**4.14** For a particle performing uniform circular motion, choose the correct statement(s) from the following:

- (a) Magnitude of particle velocity (speed) remains constant.
- (b) Particle velocity remains directed perpendicular to radius vector.
- (c) Direction of acceleration keeps changing as particle moves.
- (d) Angular momentum is constant in magnitude but direction keeps changing.

**4.15** For two vectors  $\mathbf{A}$  and  $\mathbf{B}$ ,  $|\mathbf{A} + \mathbf{B}| = |\mathbf{A} - \mathbf{B}|$  is always true when

- (a)  $|\mathbf{A}| = |\mathbf{B}| \neq 0$
- (b)  $\mathbf{A} \perp \mathbf{B}$
- (c)  $|\mathbf{A}| = |\mathbf{B}| \neq 0$  and  $\mathbf{A}$  and  $\mathbf{B}$  are parallel or anti parallel
- (d) when either  $|\mathbf{A}|$  or  $|\mathbf{B}|$  is zero.

### VSA




Fig. 4.3

**4.16** A cyclist starts from centre O of a circular park of radius 1km and moves along the path OPRQO as shown Fig. 4.3. If he maintains constant speed of  $10\text{ms}^{-1}$ , what is his acceleration at point R in magnitude and direction?

**4.17** A particle is projected in air at some angle to the horizontal, moves along parabola as shown in Fig. 4.4, where  $x$  and  $y$  indicate horizontal and vertical directions, respectively. Show in the diagram, direction of velocity and acceleration at points A, B and C.




Fig. 4.4

**4.18** A ball is thrown from a roof top at an angle of  $45^\circ$  above the horizontal. It hits the ground a few seconds later. At what point during its motion, does the ball have

- (a) greatest speed.
- (b) smallest speed.
- (c) greatest acceleration?

Explain

**4.19** A football is kicked into the air vertically upwards. What is its  
(a) acceleration, and (b) velocity at the highest point?

**4.20** **A**, **B** and **C** are three non-collinear, non co-planar vectors. What can you say about direction of  $\mathbf{A} \times (\mathbf{B} \times \mathbf{C})$ ?

## SA

**4.21** A boy travelling in an open car moving on a levelled road with constant speed tosses a ball vertically up in the air and catches it back. Sketch the motion of the ball as observed by a boy standing on the footpath. Give explanation to support your diagram.

**4.22** A boy throws a ball in air at  $60^\circ$  to the horizontal along a road with a speed of  $10 \text{ m/s}$  ( $36 \text{ km/h}$ ). Another boy sitting in a passing by car observes the ball. Sketch the motion of the ball as observed by the boy in the car, if car has a speed of ( $18 \text{ km/h}$ ). Give explanation to support your diagram.

**4.23** In dealing with motion of projectile in air, we ignore effect of air resistance on motion. This gives trajectory as a parabola as you have studied. What would the trajectory look like if air resistance is included? Sketch such a trajectory and explain why you have drawn it that way.

**4.24** A fighter plane is flying horizontally at an altitude of  $1.5 \text{ km}$  with speed  $720 \text{ km/h}$ . At what angle of sight (w.r.t. horizontal) when the target is seen, should the pilot drop the bomb in order to attack the target?

**4.25** (a) Earth can be thought of as a sphere of radius  $6400 \text{ km}$ . Any object (or a person) is performing circular motion around the axis of earth due to earth's rotation (period 1 day). What is acceleration of object on the surface of the earth (at equator) towards its centre? what is it at latitude  $\theta$ ? How does these accelerations compare with  $g = 9.8 \text{ m/s}^2$ ?

- (b) Earth also moves in circular orbit around sun once every year with an orbital radius of  $1.5 \times 10^{11} m$ . What is the acceleration of earth (or any object on the surface of the earth) towards the centre of the sun? How does this acceleration compare with  $g = 9.8 \text{ m/s}^2$ ?

$$\left( \text{Hint : acceleration } \frac{V^2}{R} = \frac{4\pi^2 R}{T^2} \right)$$

- 4.26** Given below in column I are the relations between vectors **a**, **b** and **c** and in column II are the orientations of **a**, **b** and **c** in the XY plane. Match the relation in column I to correct orientations in column II.

**Column I**

**Column II**


(a)  $\mathbf{a} + \mathbf{b} = \mathbf{c}$

(i)




(b)  $\mathbf{a} - \mathbf{c} = \mathbf{b}$

(ii)



(c)  $\mathbf{b} - \mathbf{a} = \mathbf{c}$

(iii)



(d)  $\mathbf{a} + \mathbf{b} + \mathbf{c} = 0$

(iv)



**4.27** If  $|\mathbf{A}| = 2$  and  $|\mathbf{B}| = 4$ , then match the relations in column I with the angle  $\theta$  between  $\mathbf{A}$  and  $\mathbf{B}$  in column II.

<b>Column I</b>	<b>Column II</b>
(a) $\mathbf{A} \cdot \mathbf{B} = 0$	(i) $\theta = 0^\circ$
(b) $\mathbf{A} \cdot \mathbf{B} = +8$	(ii) $\theta = 90^\circ$
(c) $\mathbf{A} \cdot \mathbf{B} = 4$	(iii) $\theta = 180^\circ$
(d) $\mathbf{A} \cdot \mathbf{B} = -8$	(iv) $\theta = 60^\circ$

**4.28** If  $|\mathbf{A}| = 2$  and  $|\mathbf{B}| = 4$ , then match the relations in column I with the angle  $\theta$  between A and B in column II

<b>Column I</b>	<b>Column II</b>
(a) $ \mathbf{A} \times \mathbf{B}  = 0$	(i) $\theta = 30^\circ$
(b) $ \mathbf{A} \times \mathbf{B}  = 8$	(ii) $\theta = 45^\circ$
(c) $ \mathbf{A} \times \mathbf{B}  = 4$	(iii) $\theta = 90^\circ$
(d) $ \mathbf{A} \times \mathbf{B}  = 4\sqrt{2}$	(iv) $\theta = 0^\circ$

## LA

**4.29** A hill is 500 m high. Supplies are to be sent across the hill using a canon that can hurl packets at a speed of 125 m/s over the hill. The canon is located at a distance of 800m from the foot of hill and can be moved on the ground at a speed of 2 m/s; so that its distance from the hill can be adjusted. What is the shortest time in which a packet can reach on the ground across the hill ? Take  $g = 10 \text{ m/s}^2$ .

**4.30** A gun can fire shells with maximum speed  $v_o$  and the maximum horizontal range that can be achieved is  $R = \frac{v_o^2}{g}$ .




Fig 4.5

If a target farther away by distance  $\Delta x$  (beyond R) has to be hit with the same gun (Fig 4.5), show that it could be achieved by raising the gun to a height at least

$$h = \Delta x \left[ 1 + \frac{\Delta x}{R} \right]$$

(Hint : This problem can be approached in two different ways:

- (i) Refer to the diagram: target T is at horizontal distance  $x = R + \Delta x$  and below point of projection  $y = -h$ .
- (ii) From point P in the diagram: Projection at speed  $v_0$  at an angle  $\theta$  below horizontal with height  $h$  and horizontal range  $\Delta x$ .)

**4.31** A particle is projected in air at an angle  $\beta$  to a surface which itself is inclined at an angle  $\alpha$  to the horizontal (Fig. 4.6).

- (a) Find an expression of range on the plane surface (distance on the plane from the point of projection at which particle will hit the surface).
- (b) Time of flight.
- (c)  $\beta$  at which range will be maximum.




Fig. 4.6

(Hint : This problem can be solved in two different ways:

- (i) Point P at which particle hits the plane can be seen as intersection of its trajectory (parabola) and straight line. Remember particle is projected at an angle  $(\alpha + \beta)$  w.r.t. horizontal.
- (ii) We can take  $x$ -direction along the plane and  $y$ -direction perpendicular to the plane. In that case resolve  $\mathbf{g}$  (acceleration due to gravity) in two different components,  $g_x$  along the plane and  $g_y$  perpendicular to the plane. Now the problem can be solved as two independent motions in  $x$  and  $y$  directions respectively with time as a common parameter.)




Fig 4.7

**4.32** A particle falling vertically from a height hits a plane surface inclined to horizontal at an angle  $\theta$  with speed  $v_0$  and rebounds elastically (Fig 4.7). Find the distance along the plane where it will hit second time.

(Hint: (i) After rebound, particle still has speed  $V_o$  to start.

(ii) Work out angle particle speed has with horizontal after it rebounds.

(iii) Rest is similar to if particle is projected up the incline.)

- 4.33** A girl riding a bicycle with a speed of 5 m/s towards north direction, observes rain falling vertically down. If she increases her speed to 10 m/s, rain appears to meet her at  $45^\circ$  to the vertical. What is the speed of the rain? In what direction does rain fall as observed by a ground based observer?

(Hint: Assume north to be  $\hat{\mathbf{i}}$  direction and vertically downward to be  $-\hat{\mathbf{j}}$ . Let the rain velocity  $\mathbf{v}_r$  be  $a\hat{\mathbf{i}} + b\hat{\mathbf{j}}$ . The velocity of rain as observed by the girl is always  $\mathbf{v}_r - \mathbf{v}_{girl}$ . Draw the vector diagram/s for the information given and find  $a$  and  $b$ . You may draw all vectors in the reference frame of ground based observer.)

- 4.34** A river is flowing due east with a speed 3m/s. A swimmer can swim in still water at a speed of 4 m/s (Fig. 4.8).

- If swimmer starts swimming due north, what will be his resultant velocity (magnitude and direction)?
- If he wants to start from point A on south bank and reach opposite point B on north bank,
  - which direction should he swim?
  - what will be his resultant speed?
- From two different cases as mentioned in (a) and (b) above, in which case will he reach opposite bank in shorter time?




Fig. 4.8

- 4.35** A cricket fielder can throw the cricket ball with a speed  $v_o$ . If he throws the ball while running with speed  $u$  at an angle  $\theta$  to the horizontal, find

- the effective angle to the horizontal at which the ball is projected in air as seen by a spectator.
- what will be time of flight?
- what is the distance (horizontal range) from the point of projection at which the ball will land?

(d) find  $\theta$  at which he should throw the ball that would maximise the horizontal range as found in (iii).

(e) how does  $\theta$  for maximum range change if  $u > v_o$ ,  $u = v_o$ ,  $u < v_o$ ?

(f) how does  $\theta$  in (v) compare with that for  $u = 0$  (i.e.  $45^\circ$ )?

**4.36** Motion in two dimensions, in a plane can be studied by expressing position, velocity and acceleration as vectors in Cartesian co-ordinates  $\mathbf{A} = A_x \hat{\mathbf{i}} + A_y \hat{\mathbf{j}}$  where  $\hat{\mathbf{i}}$  and  $\hat{\mathbf{j}}$  are unit vector along  $x$  and  $y$  directions, respectively and  $A_x$  and  $A_y$  are corresponding components of  $\mathbf{A}$  (Fig. 4.9). Motion can also be studied by expressing vectors in circular polar co-ordinates as  $\mathbf{A} = A_r \hat{\mathbf{r}} + A_\theta \hat{\boldsymbol{\theta}}$

where  $\hat{\mathbf{r}} = \frac{\mathbf{r}}{r} = \cos \theta \hat{\mathbf{i}} + \sin \theta \hat{\mathbf{j}}$  and  $\hat{\boldsymbol{\theta}} = -\sin \theta \hat{\mathbf{i}} + \cos \theta \hat{\mathbf{j}}$  are unit vectors along direction in which 'r' and ' $\theta$ ' are increasing.




Fig. 4.9

(a) Express  $\hat{\mathbf{i}}$  and  $\hat{\mathbf{j}}$  in terms of  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$ .

(b) Show that both  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$  are unit vectors and are perpendicular to each other.

(c) Show that  $\frac{d}{dt}(\hat{\mathbf{r}}) = \omega \hat{\boldsymbol{\theta}}$  where  $\omega = \frac{d\theta}{dt}$  and  $\frac{d}{dt}(\hat{\boldsymbol{\theta}}) = -\omega \hat{\mathbf{r}}$

(d) For a particle moving along a spiral given by  $\mathbf{r} = a\theta \hat{\mathbf{r}}$ , where  $a = 1$  (unit), find dimensions of 'a'.

(e) Find velocity and acceleration in polar vector representation for particle moving along spiral described in (d) above.




Fig. 4.10

**4.37** A man wants to reach from A to the opposite corner of the square C (Fig. 4.10). The sides of the square are 100 m. A central square of 50m  $\times$  50m is filled with sand. Outside this square, he can walk at a speed 1 m/s. In the central square, he can walk only at a speed of  $v$  m/s ( $v < 1$ ). What is smallest value of  $v$  for which he can reach faster via a straight path through the sand than any path in the square outside the sand?